

JEE Main 2020 Paper




Date : 2nd September 2020

Time : 09 : 00 am - 12 : 00 pm

Subject : Chemistry

1. The increasing order of the following compounds towards HCN addition is:




- (1) (iii) < (i) < (iv) < (ii) (2) (iii) < (iv) < (i) < (ii)
 (3) (i) < (iii) < (iv) < (ii) (4) (iii) < (iv) < (ii) < (i)

Sol.

1

In HCN, CN⁻ acts as nucleophile, attack first that -CHO group which has maximum positive charge. The magnitude of the (+ve) charge increases by -M and -I group. So reactivity order will be



So, option (1) is correct answer.

2. Which of the following is used for the preparation of colloids?

- (1) Van Arkel Method (2) Ostwald Process
 (3) Mond Process (4) Bredig's Arc Method

Sol.

4


Bredig's Arc method
 Chapter name surface chemistry

3. An open beaker of water in equilibrium with water vapour is in a sealed container. When a few grams of glucose are added to the beaker of water, the rate at which water molecules:



- (1) leaves the vapour increases (2) leaves the solution increases
 (3) leaves the vapour decreases (4) leaves the solution decreases

Sol.


1



Vap. press = P⁰



Vap. press = P_s



Backward shift
 vapours ↓

Hence Rate at which water molecules leaves the vap. increases.

JEE Main 2020 Paper



4. For octahedral Mn(II) and tetrahedral Ni(II) complexes, consider the following statements:

- (I) both the complexes can be high spin.
- (II) Ni(II) complex can very rarely be low spin.
- (III) with strong field ligands, Mn(II) complexes can be low spin.
- (IV) aqueous solution of Mn(II) ions is yellow in colour.

The correct statements are:

- (1) (I), (III) and (IV) only
- (2) (I), (II) and (III) only
- (3) (II), (III) and (IV) only
- (4) (I) and (II) only


Sol. 2

Mn^{2+} $[Ar]3d^5$ it can form low spin as well as high spin complex depending upon nature of ligand same of Ni^{2+} ion with coordination no 4. It can be dsp^2 or sp^3 i.e low spin or high spin depending open nature of ligand.

5. The statement that is not true about ozone is:



- (1) in the stratosphere, it forms a protective shield against UV radiation.
- (2) in the atmosphere, it is depleted by CFCs.
- (3) in the stratosphere, CFCs release chlorine free radicals (Cl) which reacts with O_3 to give chlorine dioxide radicals.
- (4) it is a toxic gas and its reaction with NO gives NO_2 .

Sol. 3



Hence option (3)

6. Consider the following reactions:




'x', 'y' and 'z' in these reactions are respectively.

- (1) 4, 5 & 5
- (2) 5, 4 & 5
- (3) 5, 6 & 5
- (4) 4, 6 & 5



JEE Main 2020 Paper



Sol. 4



α - & β -alkyl
Glucose



$(\text{CH}_3\text{CO})_2\text{O}$ reacts with $-\text{OH}$ group to form acetyl derivative, so as the no. of $-\text{OH}$ group no. of eq. of $(\text{CH}_3\text{CO})_2\text{O}$ will be used


So, $x = 4$

$y = 6$

$z = 5$


So, option (4) will be correct answer.

7. The IUPAC name for the following compound is:



- (1) 2,5-dimethyl-5-carboxy-hex-3-enal (2) 2,5-dimethyl-6-oxo-hex-3-enoic acid
(3) 6-formyl-2-methyl-hex-3-enoic acid (4) 2,5-dimethyl-6-carboxy-hex-3-enal

Sol. 2



2,5-Dimethyl-6-oxohex-3-enoic acid

JEE Main 2020 Paper



8. For the following Assertion and Reason, the correct option is

Assertion (A): When Cu (II) and sulphide ions are mixed, they react together extremely quickly to give a solid.

Reason (R): The equilibrium constant of $\text{Cu}^{2+}(\text{aq}) + \text{S}^{2-}(\text{aq}) \rightleftharpoons \text{CuS}(\text{s})$ is high because the solubility product is low.

(1) **(A)** is false and **(R)** is true.

(2) Both **(A)** and **(R)** are false.

(3) Both **(A)** and **(R)** are true but **(R)** is not the explanation for **(A)**.

(4) Both **(A)** and **(R)** are true but **(R)** is the explanation for **(A)**.




Sol. 4

(A) is (B) true &

(R) is correct explanation of (A)

Ans. 4

9. Which one of the following graphs is not correct for ideal gas?



d = Density, P = Pressure, T = Temperature

(1) I

(2) IV

(3) III


(4) II

Sol. 4


For ideal Gas

$$d = \frac{P \times M}{RT}$$


d v/s $T \rightarrow$ Hyperbolic



d v/s $\frac{1}{T} \rightarrow$ St. line



d v/s $p \rightarrow$ St line



\therefore 'II' Graph is incorrect
Ans (4)

JEE Main 2020 Paper



10. While titrating dilute HCl solution with aqueous NaOH, which of the following will not be required?



- (1) Bunsen burner and measuring cylinder (2) Burette and porcelain tile
(3) Clamp and phenolphthalein (4) Pipette and distilled water

Sol. 1

Bunsen Burner & measuring cylinder are not Required. As titration is already an exothermic process

Ans.(1)

11. In Carius method of estimation of halogen, 0.172 g of an organic compound showed presence of 0.08 g of bromine. Which of these is the correct structure of the compound?

- (1) $\text{H}_3\text{C}-\text{Br}$ (2)  (3)  (4) $\text{H}_3\text{C}-\text{CH}_2-\text{Br}$

Sol. 3

Carius method

$$\text{mass \% of 'Br'} = \frac{0.08}{0.172} \times 100 = \frac{8000}{172} = 46.51\%$$

$$\text{option (1) mass \%} = \frac{80}{95} \times 100$$

$$\text{(2) mass \%} = \frac{2 \times 80 \times 100}{252}$$

$$\text{(3) mass \%} = \frac{1 \times 80 \times 100}{80 + 72 + 6 + 14} = \frac{8000}{172} \%$$

$$\text{(4) mass \%} = \frac{1 \times 80 \times 100}{109} \%$$

Option (3) matches with the given mass percentage value


Ans (3)

12. On heating compound (A) gives a gas (B) which is a constituent of air. This gas when treated with H_2 in the presence of a catalyst gives another gas (C) which is basic in nature. (A) should not be:





- (1) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ (2) NaN_3 (3) NH_4NO_2 (4) $\text{Pb}(\text{NO}_3)_2$

Sol. 4

The gas (B) is N_2 which is found in air




(Basic in nature)




JEE Main 2020 Paper



13. The major product in the following reaction is:



Sol. 3



Option (3) is correct answer.





14. In general, the property (magnitudes only) that shows an opposite trend in comparison to other properties across a period is:

- (1) Ionization enthalpy (2) Electronegativity
 (3) Atomic radius (4) Electron gain enthalpy

Sol. 3

Ionisation energy, electronegativity & electron gain enthalpy increase across a period but atomic radius decreases

15. The figure that is not a direct manifestation of the quantum nature of atoms is:




JEE Main 2020 Paper




Sol. 4

Internal energy of 'Ar' or any gas, has nothing to do with Quantum nature of atom hence




Ans. option (4)

16. The major aromatic product C in the following reaction sequence will be :



Sol. 3



Option (3) is correct answer.

JEE Main 2020 Paper



- 20.** The metal mainly used in devising photoelectric cells is:
 (1) Li (2) Cs (3) Rb (4) Na

Sol. 2
 'Cs' is used in photoelectric cell as its ionisation energy is lowest
 Hence Ans (2)

- 21.** The mass of gas adsorbed, x , per unit mass of adsorbate, m , was measured at various pressures, p . A graph between $\log \frac{x}{m}$ and $\log p$ gives a straight line with slope equal to 2 and the intercept equal to 0.4771. The value of $\frac{x}{m}$ at a pressure of 4 atm is: (Given $\log 3 = 0.4771$)

Sol. 48
 $\frac{x}{m} = kP^{1/n}$
 $\log (x / m) = \log_{(k)} + \frac{1}{n} \log(p)$
 $y = c + mx$
 Intercept C = $\log_k = 0.4771$
 slop = $\frac{1}{n} = 2, k = 3$
 $\frac{x}{m} = k(P)^{1/n}$ at P = 4 atm
 $= 3(4)^2$
 $\frac{x}{m} = 3 \times 16 = 48$ Ans

- 22.** The Gibbs energy change (in J) for the given reaction at $[Cu^{2+}] = [Sn^{2+}] = 1$ M and 298 K is:
 $Cu(s) + Sn^{2+}(aq.) \rightarrow Cu^{2+}(aq.) + Sn(s)$

($E_{Sn^{2+}|Sn}^{\circ} = -0.16V, E_{Cu^{2+}|Cu}^{\circ} = 0.34V$, Take $F = 96500$ C mol⁻¹)

Sol. 96500
 $Cu(s) + Sn^{2+}(aq) \rightleftharpoons Cu^{2+}(aq) + Sn(s)$
 $E_{cell}^{\circ} = -0.16 - 0.34$
 $= -0.50$
 $\Delta G^{\circ} = -nF E_{cell}^{\circ}$
 $= -2 \times 96500 \times (-0.5)$
 $= +96500$
 $\Delta G = \Delta G^{\circ} + RT \ln Q$
 $= 96500 + \frac{25}{3} \times 298 \times 2.303 \log (1)$
 $\Delta G = 96500$ Joules


- 23.** The internal energy change (in J) when 90 g of water undergoes complete evaporation at 100° C is _____.
 (Given: ΔH_{vap} for water at 373 K = 41 kJ/mol, $R = 8.314$ JK⁻¹ mol⁻¹)

Sol. $H_2O(l) \rightarrow H_2O(g)$
 $\Delta E_{vap} = \Delta H_{vap} - \Delta ngRT$
 $= 41000 \times 5 - 5 \times 8.314 \times 373$
 $= 189494.39$




JEE Main 2020 Paper




24. The oxidation states of iron atoms in compounds (A), (B) and (C), respectively, are x, y and z. The sum of x, y and z is _____.




Sol. 6



25. The number of chiral carbons present in the molecule given below is _____.



Sol. 5



Total chiral carbon = 5